

Furthermore, perform the summations over all spin states and the polarizations, with which you get

$$\overline{|\mathcal{M}|^2} = \frac{1}{3} \sum_{s,\lambda} |\mathcal{M}|^2 = \frac{g^4 m_W^2}{3 [(p+q)^2 - m_W^2]^2} \left\{ (p \cdot q) + \frac{2(p \cdot k)(q \cdot k)}{m_W^2} \right\} \quad (34)$$

For the calculation of the decay width use the convention

$$(k-p)^2 = t, \quad (k-q)^2 = u, \quad (k-l)^2 = s, \quad (35)$$


and the formula for the three-particle phase space,

$$d\text{Lips} = \frac{1}{128\pi^3 m_W^2} \int ds dt. \quad (36)$$

Consider the kinematic limits of the integration variables based on the results of example 15 (WS) and give the result for the differential width $\frac{d^2\Gamma}{ds dt}$. Furthermore make the assumption $m_H \ll m_W$ (we know, experimentally $m_H > m_W$) and show that in this approximation the result for $d\Gamma/ds$ is

$$\frac{d\Gamma}{ds} = \frac{g^4}{768\pi^3 m_W} \left(-\frac{1}{12} + \frac{s}{12 m_W^2} + \frac{s}{s - m_W^2} \right). \quad (37)$$

14. We have the reaction $e^- + \nu_\mu \rightarrow e^- + \nu_\mu$.



Extract the necessary Feynman rules for the vertices from the Lagrangian density

$$\mathcal{L}_{NC} = \frac{g}{\cos \theta_W} J_\mu^{NC} Z^\mu, \quad (38)$$

where

$$J_\mu^{NC} = \frac{1}{2} \left[g_L^f \bar{f} \gamma_\mu (1 - \gamma_5) f + g_R^f \bar{f} \gamma_\mu (1 + \gamma_5) f \right] \quad (39)$$

and

$$g_L^f = T_3(f_L) - Q(f_L) \sin^2 \theta_W, \quad (40)$$

$$g_R^f = T_3(f_R) - Q(f_R) \sin^2 \theta_W. \quad (41)$$

The Feynman rule for the Z-propagator in the unitary gauge can be found in the lecture. Calculate the matrix element, which is

$$\mathcal{M} = \frac{ig^2}{4\cos^2\theta_W} \frac{g_L^\nu}{q^2 - m_Z^2} \bar{u}(k')\gamma_\mu(1-\gamma_5)u(k) (\bar{u}(p')\gamma^\mu (g_L^e(1-\gamma_5) + g_R^e(1+\gamma_5)) u(p)) . \quad (42)$$

Further calculate the cross section $d\sigma/d\cos\theta_{CMS}$ of this reaction. The result is

$$\frac{d\sigma}{d\cos\theta_{CMS}}(e^-\nu_\mu \rightarrow e^-\nu_\mu) = \frac{g^4}{128\pi\cos^4\theta_W} \frac{(g_R^e)^2 s^2 + (g_L^e)^2 u^2}{s(t-m_Z^2)^2}, \quad (43)$$

with the Mandelstam variables $s = (p+k)^2$, $t = (k-k')^2$, and $u = (k-p')^2$. What is the result for the reaction $e^- + \bar{\nu}_\mu \rightarrow e^- + \bar{\nu}_\mu$?

15. Calculate the differential cross section of the reaction $e^+ + e^- \rightarrow b + \bar{b}$ at the Z^0 pole, i.e. the four-momentum transmitted to Z^0 is $|q| \sim m_{Z^0}$. The mass of the bottom quark is not neglected. Take the Feynman rules from the lecture notes.

First show that the matrix element \mathcal{M} has the form

$$\mathcal{M} = \frac{ig^2}{4\cos^2\theta_W} \frac{1}{q^2 - m_Z^2} \bar{v}(p_2)\gamma_\mu(v_e + a_e\gamma_5)u(p_1)\bar{u}(p_3)\gamma^\mu(v_b + a_b\gamma_5)v(p_4), \quad (44)$$

and the result has the form

$$\frac{d\sigma}{d\cos\theta} = \frac{1}{16\pi s} \frac{g^4}{(s - m_Z^2)^2} \left[2Am_b^2 s + (B+C)(s+t-m_b^2)^2 + (B-C)(m_b^2-t)^2 \right], \quad (45)$$

with the factors $A = \frac{1}{16\cos^4\theta_W}(v_b^2 - a_b^2)(v_e^2 + a_e^2)$, $B = \frac{1}{16\cos^4\theta_W}(v_b^2 + a_b^2)(v_e^2 + a_e^2)$ and $C = \frac{1}{4\cos^4\theta_W}v_b a_b v_e a_e$. The Mandelstam variable s is the square of the momentum of the exchanged Z boson and t is the square of the difference of the momenta of the incoming electron and outgoing b quark. Sketch also how to obtain the total cross section.

