

# Complexity Theory

VU 181.142, SS 2014

Homework Assignment 2

Name: N.N.  
Matr-Nr: xxxxxxxx  
Begin: 25 March, 2014  
Submission Deadline: 8 April, 2014  
send to: complexity@dbai.tuwien.ac.at  
Maximum credits: 10

**Exercise 1 (5 credits)** Recall the problem reduction from an arbitrary language  $L \in \text{NP}$  to the **SAT**-problem given in the lecture. In particular, recall the construction of an instance  $R(x)$  of **SAT** from an arbitrary instance  $x$  of  $L$ . Give a rigorous proof of the correctness of this reduction, i.e.  $x \in L \Leftrightarrow R(x) \in \text{SAT}$ .

**Hint.** Prove both directions of the equivalence separately. The intended meaning of the propositional atoms in  $R(x)$  is clear. You have to be careful, what is given, what is constructed (or defined), and what has to be proved.

- Suppose that  $x \in L$ .
  - given: Then we know that there exists a successful computation of the NTM  $T$  on input  $x$ . By our assumption, this computation consists of exactly  $N$  steps. Let  $conf_0, \dots, conf_N$  denote the configurations of the NTM  $T$  along this computation.
  - constructed/defined: We define a truth assignment  $\mathcal{I}$  appropriate to  $R(x)$  according to the intended meaning of the propositional atoms in  $R(x)$ .
  - to be proved: It remains to show that all conjuncts in  $R(x)$  are indeed satisfied by  $\mathcal{I}$ . For this purpose, you have to inspect all groups of conjuncts in  $R(x)$  and argue that each of them is true in  $\mathcal{I}$ .
- Suppose that  $R(x) \in \text{SAT}$ .
  - given: Then there exists a satisfying truth assignment  $\mathcal{I}$  of  $R(x)$ .
  - constructed/defined: We can construct a sequence  $conf_0, \dots, conf_N$  of configurations of the NTM  $T$  according to the intended meaning of the propositional atoms in  $R(x)$ .
  - to be proved: First argue that the configurations  $conf_0, \dots, conf_N$  are well-defined (by using the the fact that  $\mathcal{I}$  satisfies all conjuncts of  $R(x)$ ). It remains to show that there exists a computation of  $T$  on input  $x$  which produces exactly this sequence of configurations  $conf_0, \dots, conf_N$ . Note that,

in particular, this is a successful computation by the conjunct  $state_{s_m}[N]$  (with  $s_m = \text{“yes”}$ ) in  $R(x)$  and by the intended meaning of  $state_{s_m}[N]$ . For this purpose, you have to show by induction on  $\tau$  that there exists a computation of  $T$  on input  $x$  whose first  $\tau$  configurations are  $conf_0, \dots, conf_\tau$ .

**For your convenience.** The groups of conjuncts in  $R(x)$  are recalled below.

**1. Initialization facts.**

$symbol_{\triangleright}[0, 0]$   
 $symbol_{\sigma}[0, \pi]$  for  $1 \leq \pi \leq |x|$ , where  $x_\pi = \sigma$   
 $symbol_{\sqcup}[0, \pi]$  for  $|x| < \pi \leq N$   
 $cursor[0, 0]$   
 $state_{s_0}[0]$

**2. Transition rules.** For each pair  $(s, \sigma)$  of state  $s$  and symbol  $\sigma$  let  $\langle s, \sigma, s'_1, \sigma'_1, d_1 \rangle, \dots, \langle s, \sigma, s'_k, \sigma'_k, d_k \rangle$  denote all possible transitions according to the transition relation  $\Delta$  (for the cursor movements, we write  $d_i \in \{-1, 0, 1\}$  rather than  $d_i \in \{\leftarrow, -, \rightarrow\}$ ).

Then  $R(x)$  contains the following conjuncts for each value of  $\tau$  and  $\pi$  such that  $0 \leq \tau < N$  and  $0 \leq \pi < N$

$state_s[\tau] \wedge symbol_{\sigma}[\tau, \pi] \wedge cursor[\tau, \pi] \rightarrow$   
 $[(state_{s'_1}[\tau + 1] \wedge symbol_{\sigma'_1}[\tau + 1, \pi] \wedge cursor[\tau + 1, \pi + d_1]) \vee \dots \vee$   
 $(state_{s'_k}[\tau + 1] \wedge symbol_{\sigma'_k}[\tau + 1, \pi] \wedge cursor[\tau + 1, \pi + d_k])]$

**3. Uniqueness constraints.** Let  $K = \{s_0, \dots, s_m\}$  and  $\Sigma = \{\sigma_1, \dots, \sigma_n\}$ .

Then  $R(x)$  contains the following formulae for each value of  $\tau$  and  $\pi$  such that  $0 \leq \tau \leq N$ ,  $0 \leq \pi \leq N$ ,  $0 \leq i \leq m$ , and  $1 \leq j \leq n$ .

$state_{s_i}[\tau] \leftrightarrow (\neg state_{s_0}[\tau] \wedge \dots \wedge \neg state_{s_{i-1}}[\tau] \wedge$   
 $\wedge \neg state_{s_{i+1}}[\tau] \wedge \dots \wedge \neg state_{s_m}[\tau])$   
 $cursor[\tau, \pi] \leftrightarrow (\neg cursor[\tau, 0] \wedge \dots \wedge \neg cursor[\tau, \pi - 1] \wedge$   
 $\wedge \neg cursor[\tau, \pi + 1] \wedge \dots \wedge \neg cursor[\tau, N])$   
 $symbol_{\sigma_j}[\tau, \pi] \leftrightarrow (\neg symbol_{\sigma_1}[\tau, \pi] \wedge \dots \wedge \neg symbol_{\sigma_{j-1}}[\tau, \pi] \wedge$   
 $\wedge \neg symbol_{\sigma_{j+1}}[\tau, \pi] \wedge \dots \wedge \neg symbol_{\sigma_n}[\tau, \pi])$

**4. Inertia rules.**  $R(x)$  contains the following conjuncts for each value  $\tau, \pi, \pi', \sigma$ , where  $0 \leq \tau < N$ ,  $0 \leq \pi < \pi' \leq N$ , and  $\sigma \in \Sigma$ ,

$symbol_{\sigma}[\tau, \pi], cursor[\tau, \pi'] \rightarrow symbol_{\sigma}[\tau + 1, \pi]$   
 $symbol_{\sigma}[\tau, \pi'], cursor[\tau, \pi] \rightarrow symbol_{\sigma}[\tau + 1, \pi']$

**5. Acceptance.** Let  $s_m = \text{“yes”}$ .

Then  $R(x)$  contains the following atom as a conjunct:

$state_{s_m}[N]$ .

**Exercise 2 (5 credits)** Recall the basic, polynomial-time decision procedure for HORN-SAT (see cc04.pdf). The correctness of this decision procedure relies on the following lemma:

**Lemma.** *Let  $\varphi$  be a propositional Horn-formula. Let  $Y$  denote the set of atoms which are obtained by initializing  $Y$  to the set of facts in  $\varphi$  and by exhaustively applying the rules in  $\varphi$  to  $Y$ . Then, for every atom  $x$  in  $\varphi$ , the following equivalence holds:  $x \in Y \Leftrightarrow x$  is implied by the facts and rules in  $\varphi$ .*

Give a rigorous proof of this lemma.