

Complexity Theory

VU 181.142, SS 2015

Homework Assignment 2

Name: N.N.
Matr-Nr: xxxxxxxx
Begin: 24 March, 2015
Submission Deadline: 21 April, 2015
send to: complexity@dbai.tuwien.ac.at
Maximum credits: 10

Exercise 1 (5 credits) Recall the problem reduction from an arbitrary language $L \in \text{NP}$ to the **SAT**-problem given in the lecture. In particular, recall the construction of an instance $R(x)$ of **SAT** from an arbitrary instance x of L . Give a rigorous proof of the correctness of this reduction, i.e. $x \in L \Leftrightarrow R(x) \in \text{SAT}$.

Hint. Prove both directions of the equivalence separately. The intended meaning of the propositional atoms in $R(x)$ is clear. You have to be careful, what is given, what is constructed (or defined), and what has to be proved.

- Suppose that $x \in L$.
 - given: Then we know that there exists a successful computation of the NTM T on input x . By our assumption, this computation consists of exactly N steps. Let $conf_0, \dots, conf_N$ denote the configurations of the NTM T along this computation.
 - constructed/defined: We define a truth assignment \mathcal{I} appropriate to $R(x)$ according to the intended meaning of the propositional atoms in $R(x)$.
 - to be proved: It remains to show that all conjuncts in $R(x)$ are indeed satisfied by \mathcal{I} . For this purpose, you have to inspect all groups of conjuncts in $R(x)$ and argue that each of them is true in \mathcal{I} .
- Suppose that $R(x) \in \text{SAT}$.
 - given: Then there exists a satisfying truth assignment \mathcal{I} of $R(x)$.
 - constructed/defined: We can construct a sequence $conf_0, \dots, conf_N$ of configurations of the NTM T according to the intended meaning of the propositional atoms in $R(x)$.
 - to be proved: First argue that the configurations $conf_0, \dots, conf_N$ are well-defined (by using the the fact that \mathcal{I} satisfies all conjuncts of $R(x)$). It remains to show that there exists a computation of T on input x which produces exactly this sequence of configurations $conf_0, \dots, conf_N$. Note that,

in particular, this is a successful computation by the conjunct $state_{s_m}[N]$ (with $s_m = \text{“yes”}$) in $R(x)$ and by the intended meaning of $state_{s_m}[N]$. For this purpose, you have to show by induction on τ that there exists a computation of T on input x whose first τ configurations are $conf_0, \dots, conf_\tau$.

For your convenience. The groups of conjuncts in $R(x)$ are recalled below.

1. Initialization facts.

$symbol_{\triangleright}[0, 0]$
 $symbol_{\sigma}[0, \pi]$ for $1 \leq \pi \leq |x|$, where $x_\pi = \sigma$
 $symbol_{\sqcup}[0, \pi]$ for $|x| < \pi \leq N$
 $cursor[0, 0]$
 $state_{s_0}[0]$

2. Transition rules. For each pair (s, σ) of state s and symbol σ let $\langle s, \sigma, s'_1, \sigma'_1, d_1 \rangle, \dots, \langle s, \sigma, s'_k, \sigma'_k, d_k \rangle$ denote all possible transitions according to the transition relation Δ (for the cursor movements, we write $d_i \in \{-1, 0, 1\}$ rather than $d_i \in \{\leftarrow, -, \rightarrow\}$).

Then $R(x)$ contains the following conjuncts for each value of τ and π such that $0 \leq \tau < N$ and $0 \leq \pi < N$

$state_s[\tau] \wedge symbol_{\sigma}[\tau, \pi] \wedge cursor[\tau, \pi] \rightarrow$
 $[(state_{s'_1}[\tau + 1] \wedge symbol_{\sigma'_1}[\tau + 1, \pi] \wedge cursor[\tau + 1, \pi + d_1]) \vee \dots \vee$
 $(state_{s'_k}[\tau + 1] \wedge symbol_{\sigma'_k}[\tau + 1, \pi] \wedge cursor[\tau + 1, \pi + d_k])]$

3. Uniqueness constraints. Let $K = \{s_0, \dots, s_m\}$ and $\Sigma = \{\sigma_1, \dots, \sigma_n\}$.

Then $R(x)$ contains the following formulae for each value of τ and π such that $0 \leq \tau \leq N$, $0 \leq \pi \leq N$, $0 \leq i \leq m$, and $1 \leq j \leq n$.

$state_{s_i}[\tau] \leftrightarrow (\neg state_{s_0}[\tau] \wedge \dots \wedge \neg state_{s_{i-1}}[\tau] \wedge$
 $\wedge \neg state_{s_{i+1}}[\tau] \wedge \dots \wedge \neg state_{s_m}[\tau])$
 $cursor[\tau, \pi] \leftrightarrow (\neg cursor[\tau, 0] \wedge \dots \wedge \neg cursor[\tau, \pi - 1] \wedge$
 $\wedge \neg cursor[\tau, \pi + 1] \wedge \dots \wedge \neg cursor[\tau, N])$
 $symbol_{\sigma_j}[\tau, \pi] \leftrightarrow (\neg symbol_{\sigma_1}[\tau, \pi] \wedge \dots \wedge \neg symbol_{\sigma_{j-1}}[\tau, \pi] \wedge$
 $\wedge \neg symbol_{\sigma_{j+1}}[\tau, \pi] \wedge \dots \wedge \neg symbol_{\sigma_n}[\tau, \pi])$

4. Inertia rules. $R(x)$ contains the following conjuncts for each value τ, π, π', σ , where $0 \leq \tau < N$, $0 \leq \pi < \pi' \leq N$, and $\sigma \in \Sigma$,

$symbol_{\sigma}[\tau, \pi], cursor[\tau, \pi'] \rightarrow symbol_{\sigma}[\tau + 1, \pi]$
 $symbol_{\sigma}[\tau, \pi'], cursor[\tau, \pi] \rightarrow symbol_{\sigma}[\tau + 1, \pi']$

5. Acceptance. Let $s_m = \text{“yes”}$.

Then $R(x)$ contains the following atom as a conjunct:

$state_{s_m}[N]$.

Exercise 2 (5 credits) Recall the basic, polynomial-time decision procedure for HORN-SAT (see cc04.pdf). The correctness of this decision procedure relies on the following lemma:

Lemma. *Let φ be a propositional Horn-formula. Let Y denote the set of atoms which are obtained by initializing Y to the set of facts in φ and by exhaustively applying the rules in φ to Y . Then, for every atom x in φ , the following equivalence holds: $x \in Y \Leftrightarrow x$ is implied by the facts and rules in φ .*

Give a rigorous proof of this lemma.

Terminology. We say that a formula β is implied by a formula α , if $\models (\alpha \rightarrow \beta)$ holds, i.e., every model of α is a model of β . In the above lemma, let ψ denote the conjunction consisting of the rules and facts (but not the goals) in φ . The lemma thus claims that $x \in Y \Leftrightarrow \models (\psi \rightarrow x)$ holds.